




The future Cold QCD program with the sPHENIX detector

Desmond Shangase (University of Michigan) on behalf of the sPHENIX Collaboration
RHIC/AGS Annual Users Meeting - October 22nd 2020




Contents

- sPHENIX Detector Design + Run
- Cold QCD Measurements
 - Transverse Spin Measurements
 - Unpolarized Measurements



sPHENIX Detector

- Full azimuthal detector (Central Barrel)
- Data collection expected to begin 2023
- Cold QCD Physics Program
 - Parton Dynamics (TMD PDFs)
 - Proton/Nuclear Structure (PDFs)
 - Hadronization (FFs, \hat{q} , etc.)



6

sPHENIX Detector

- Full azimuthal detector (Central Barrel)
- Data collection expected to begin 2023
- Cold QCD Physics Program
 - Parton Dynamics (TMD PDFs)
 - Proton/Nuclear Structure (PDFs)
 - Hadronization (FFs, \hat{q} , etc.)

sPH-TRG-2020-001

Year	Species	$\sqrt{s_{NN}}$ [GeV]	Cryo Weeks	Physics Weeks	Rec. Lum. $ z < 10$ cm	Samp. Lum. $ z < 10$ cm
2023	Au+Au	200	24 (28)	9 (13)	3.7 (5.7) nb ⁻¹	4.5 (6.9) nb ⁻¹
2024	$p^\dagger p^\dagger$	200	24 (28)	12 (16)	0.3 (0.4) pb ⁻¹ [5 kHz] 4.5 (6.2) pb ⁻¹ [10%-str]	45 (62) pb ⁻¹
2024	$p^\dagger + \text{Au}$	200	-	5	0.003 pb ⁻¹ [5 kHz] 0.01 pb ⁻¹ [10%-str]	0.11 pb ⁻¹
2025	Au+Au	200	24 (28)	20.5 (24.5)	13 (15) nb ⁻¹	21 (25) nb ⁻¹

Outer HCAL
gnet
ICAL

mediate Tracker
/erTeX Detector

cryo
chimney


sPHENIX Detector

- Full azimuthal detector (Central Barrel)
- Data collection expected to begin 2023
- Cold QCD Physics Program
 - Parton Dynamics (TMD PDFs)
 - Proton/Nuclear Structure (PDFs)
 - Hadronization (FFs, \hat{q} , etc.)

sPH-TRG-2020-001

Year	Species	$\sqrt{s_{NN}}$ [GeV]	Cryo Weeks	Physics Weeks	Rec. Lum. z < 10 cm	Samp. Lum. z < 10 cm
2023	Au+Au	200	24 (28)	9 (13)	3.7 (5.7) nb ⁻¹	4.5 (6.9) nb ⁻¹
2024	$p^\uparrow p^\uparrow$	200	24 (28)	12 (16)	0.3 (0.4) pb ⁻¹ [5 kHz] 4.5 (6.2) pb ⁻¹ [10%-str]	45 (62) pb ⁻¹
2024	p^\uparrow +Au	200	-	5	0.003 pb ⁻¹ [5 kHz] 0.01 pb ⁻¹ [10%-str]	0.11 pb ⁻¹
2025	Au+Au	200	24 (28)	20.5 (24.5)	13 (15) nb ⁻¹	21 (25) nb ⁻¹

Last polarized hadron collision data to come out of RHIC



sPHENIX Detector

- Full azimuthal detector (Central Barrel)
- Data collection expected to begin 2023
- Cold QCD Physics Program
 - Parton Dynamics (TMD PDFs)
 - Proton/Nuclear Structure (PDFs)
 - Hadronization (FFs, \hat{q} , etc.)

Tamamushi, S. (2017)

Nucleon Polarization

Quark Polarization

	Unpolarized (U)	Longitudinal (L)	Transverse (T)
U	Number Density f_1 		Sivers f_{1T}^\perp
L		Helicity g_{1L} 	Worm-Gear-1 g_{1T}
T	Boer-Mulders $h_{1\perp}^\perp$ 	Worm-Gear-2 $h_{1\perp}^\perp$ 	Transversity h_{1T}
			Pretzelicity h_{1T}^\perp

er HCAL
Magnet
r HCAL
CAL

mediate Tracker
S VerTeX Detector

cryo
chimney

sPHENIX Detector

- Full azimuthal detector (Central Barrel)
- Data collection expected to begin 2023
- Cold QCD Physics Program
 - Parton Dynamics (TMD PDFs)
 - Proton/Nuclear Structure (PDFs)
 - Hadronization (FFs, \hat{q} , etc.)

Tamamushi, S. (2017)

Nucleon Polarization

Quark Polarization

	Unpolarized (U)	Longitudinal (L)	Transverse (T)
U	Number Density f_1 		Sivers f_{1T}^\perp
L		Helicity g_{1L} 	Worm-Gear-1 g_{1T}
T	Boer-Mulders $h_{1\perp}^\perp$ 	Worm-Gear-2 $h_{1\perp}^\perp$ 	Transversity h_{1T}
			Pretzelosity h_{1T}^\perp

er HCAL
Magnet
r HCAL
CAL

mediate Tracker
S VerTeX Detector

cryo chimney


Transverse Spin Measurements in $p^\uparrow + p^{(\uparrow)}$ and $p^\uparrow + \text{Au}$

Gluon Dynamics via Transverse Single Spin Asymmetry A_N

Direct Photon Asymmetry

- Will be used to constrain twist-3 trigluon correlator in transversely polarized protons
 - Related to f_{1T}^\perp of gluons in the proton

$$A_N(\phi_q) = \frac{1}{P} \frac{Y^\uparrow - R \cdot Y^\downarrow}{Y^\uparrow + R \cdot Y^\downarrow} = \frac{1}{P} \frac{L(\sigma^\uparrow(\phi_q) - R \cdot \sigma^\downarrow(\phi_q))}{L(\sigma^\uparrow(\phi_q) + R \cdot \sigma^\downarrow(\phi_q))}$$




Gluon Dynamics via Transverse Single Spin Asymmetry A_N

Direct Photon Asymmetry

- Will be used to constrain twist-3 trigluon correlator in transversely polarized protons
 - Related to f_{1T}^\perp of gluons in the proton

$$A_N(\phi_q) = \frac{1}{P} \frac{Y^\uparrow - R \cdot Y^\downarrow}{Y^\uparrow + R \cdot Y^\downarrow} = \frac{1}{P} \frac{L(\sigma^\uparrow(\phi_q) - R \cdot \sigma^\downarrow(\phi_q))}{L(\sigma^\uparrow(\phi_q) + R \cdot \sigma^\downarrow(\phi_q))}$$




Gluon Dynamics via Transverse Single Spin Asymmetry A_N

Heavy Flavor Asymmetry

- Will be used to constrain twist-3 trigluon correlator in transversely polarized protons
 - Related to f_{1T}^\perp of gluons in the proton

$$A_N(\phi_q) = \frac{1}{P} \frac{Y^\uparrow - R \cdot Y^\downarrow}{Y^\uparrow + R \cdot Y^\downarrow} = \frac{1}{P} \frac{L(\sigma^\uparrow(\phi_q) - R \cdot \sigma^\downarrow(\phi_q))}{L(\sigma^\uparrow(\phi_q) + R \cdot \sigma^\downarrow(\phi_q))}$$


- Possible due to sPHENIX streaming DAQ
 - 10% of collisions will be recorded in this triggerless configuration



Glue Dynamics via Transverse Single Spin Asymmetry A_N

Gamma-jet Asymmetry

- Gluon-induced Compton scattering
 - Constrain gluon spin distribution in polarized proton
 - sPHENIX is designed to be a jet detector due to the relevance of this and similar channels to heavy-ion physics





Parton Dynamics via A_N

Gamma-jet Asymmetry

- Gluon-induced Compton scattering
 - Constrain gluon spin distribution in polarized proton
 - sPHENIX is designed to be a jet detector due to the relevance of this and similar channels to heavy-ion physics

Dijet Asymmetry

- Charge-tagging can allow for flavor-dependent Sivers asymmetry measurement





Parton Dynamics via A_N

Gamma-jet Asymmetry

- Gluon-induced Compton scattering
 - Constrain gluon spin distribution in polarized proton
 - sPHENIX is designed to be a jet detector due to the relevance of this and similar channels to heavy-ion physics

Dijet Asymmetry

- Charge-tagging can allow for flavor-dependent Sivers asymmetry measurement




arXiv:2008.05470v2 [hep-ph]

Nuclear Effects in A_N

Charged hadron Asymmetry

- Noticeable decrease in A_N amplitude in differing collision systems
 - At forward pseudorapidity and intermediate x_F
 - Currently no consensus on this behavior




Phys. Rev. Lett. 123 (2019) 12, 122001

Nuclear Effects in A_N


Charged hadron Asymmetry

- Noticeable decrease in A_N amplitude in differing collision systems
 - At forward pseudorapidity and intermediate x_F
 - Currently no consensus on this behavior
- sPHENIX to improve statistics in this region of x_F
 - Specifically for $p^\uparrow + p^\uparrow$ and $p^\uparrow + \text{Au}$ data points
 - Finer binning is expected




Unpolarized Measurements in p+p and p+Au

Nuclear Effects in Hadronization




sPH-TRG-2020-001




- Due to sPHENIX Central Barrel and Vertex Detector
 - Direct photons and charged hadrons up to ~ 45 GeV
 - Jets up to ~ 70 GeV
- Nuclear modification of hadron-in-jet distributions possible

Nuclear Effects in Hadronization




- Due to sPHENIX Central Barrel and Vertex Detector
 - Direct photons and charged hadrons up to ~ 45 GeV
 - Jets up to ~ 70 GeV
- Nuclear modification of hadron-in-jet distributions possible
 - w.r.t. z , j_T , r , etc.



Nuclear Effects in Hadronization



- Due to sPHENIX Central Barrel and Vertex Detector
 - Direct photons and charged hadrons up to ~45 GeV
 - Jets up to ~70 GeV
- Nuclear modification of hadron-in-jet distributions possible
 - w.r.t. z , j_T , r , etc.
- Similarly, can measure transport coefficient for gamma-jet systems
 - $\langle \hat{q}L \rangle / 2 \cong \langle p_{out}^2 \rangle_{pA} - \langle p_{out}^2 \rangle_{pp}$





Constraining nPDFs



- Tension exists between nPDF models

Phys. Rev. D 100, 014004


Constraining nPDFs



- Tension exists between nPDF models
- Measurement of nuclear modifications can be used to constrain existing nPDFs
- Channels expected for simultaneous analysis
 - Drell-Yan
 - Dijet
 - Photon-jet

Phys. Rev. D 100, 014004

Constraining nPDFs



Phys. Rev. D 100, 014004

- Tension exists between nPDF models
- Measurement of nuclear modifications can be used to constrain existing nPDFs
- Channels expected for simultaneous analysis
 - Drell-Yan
 - Dijet
 - Photon-jet
- Expecting improved uncertainties in gluon and antiquark nPDFs with this method
 - Particularly in shadowing region

Further Prospects

- Sivers via inclusive jet A_N
 - Yet to be measured at central rapidity
 - Uncertainty expected on the order of 10^{-4}
 - Complementary study to be done at EIC
- Collins Fragmentation Function
 - H_1^\perp = distribution of in-jet hadron transverse momentum produced by a polarized quark
 - Provides us much needed access to transversity in protons
 - h_1 = parton transverse spin polarization in a transversely polarized proton
- Interference Fragmentation Function
 - Coupling between transversity and dihadron hadronization
 - Measured via dihadron angular distributions

Summary

- Parton dynamics and cold nuclear effects can be measured/constrained with the sPHENIX detector
- Transverse spin dependent observables grant us access to
 - Gluon dynamics via photon, photon-jet (new), heavy flavor, and dijet asymmetries
 - Quark dynamics via charge-tagging in dijet channel
 - A_N nuclear and pseudorapidity dependencies via inclusive hadron measurements
- Viable spin-independent measurements at sPHENIX will contribute to understanding of transport coefficients as well as the nuclear modification of
 - Direct photons, charged hadrons (new), and inclusive jet production
 - Heavy flavor distributions in jets
 - Gluon and antiquark PDFs via Drell-Yan, dijet, and photon-jet channels