


# The jet physics program with sPHENIX

Virginia Bailey  
for the sPHENIX Collaboration  
March 1, 2022

# sPHENIX physics program


2

Jets and jet substructure




Vary momentum,  
angular scale of  
probe

Upsilon spectroscopy




Vary size of probe

Open heavy-flavor



Vary mass of probe

Cold QCD



Study cold nuclear  
matter effects

See talk on heavy-flavor physics on Friday by Thomas Marshall

# sPHENIX detector

3

## Tracking detector:

- MAPS-based Vertex Tracker (MVTX)
- Intermediate Silicon Tracker (INTT)
- Time Projection Chamber (TPC)

## Superconducting Magnet


- 1.4T solenoid magnet

## Calorimeter:

- Electromagnetic calorimeter (EMCal)
- Inner hadronic calorimeter (inner HCal)
- Outer hadronic calorimeter (outer HCal)

## High rate DAQ and trigger systems

- 15 kHz trigger



# sPHENIX calorimeter

4


Full calorimeter covers  $2\pi$  in azimuth  
and  $|\eta| < 1.1$

## EMCal:

- Sampling calorimeter of scintillating fibers embedded in tungsten blocks
- $\Delta\eta \times \Delta\phi = 0.025 \times 0.025$  towers



**Block production  
finished at UIUC**





# sPHENIX calorimeter

5


**Full calorimeter covers  $2\pi$  in azimuth and  $|\eta| < 1.1$**

## **EMCal:**

- Sampling calorimeter of scintillating fibers embedded in tungsten blocks
- $\Delta\eta \times \Delta\phi = 0.025 \times 0.025$  towers

## **Inner and outer HCal:**

- Sampling calorimeter of scintillating tiles and steel absorber plates
- $\Delta\eta \times \Delta\phi = 0.1 \times 0.1$  towers



# sPHENIX calorimeter

6

**Full calorimeter covers  $2\pi$  in azimuth and  $|\eta| < 1.1$**


## **EMCal:**

- Sampling calorimeter of scintillating fibers embedded in tungsten blocks
- $\Delta\eta \times \Delta\phi = 0.025 \times 0.025$  towers

## **Inner and outer HCal:**

- Sampling calorimeter of scintillating tiles and steel absorber plates
- $\Delta\eta \times \Delta\phi = 0.1 \times 0.1$  towers

**Calorimeters read out with SiPMs**



# sPHENIX run plan

7

Year	Species	$\sqrt{s_{NN}}$ [GeV]	Cryo Weeks	Physics Weeks	Rec. Lum. $ z  < 10$ cm	Samp. Lum. $ z  < 10$ cm
2023	Au+Au	200	24 (28)	9 (13)	3.7 (5.7) nb <sup>-1</sup>	4.5 (6.9) nb <sup>-1</sup>
2024	$p^\uparrow p^\uparrow$	200	24 (28)	12 (16)	0.3 (0.4) pb <sup>-1</sup> [5 kHz] 4.5 (6.2) pb <sup>-1</sup> [10%-str]	45 (62) pb <sup>-1</sup>
2024	$p^\uparrow + \text{Au}$	200	–	5	0.003 pb <sup>-1</sup> [5 kHz] 0.01 pb <sup>-1</sup> [10%-str]	0.11 pb <sup>-1</sup>
2025	Au+Au	200	24 (28)	20.5 (24.5)	13 (15) nb <sup>-1</sup>	21 (25) nb <sup>-1</sup>


< 1 year until data

Large luminosity in first year

# Why jet measurements at RHIC?

8



- Different QGP:
  - Temperature/temperature evolution different between LHC and RHIC



# Why jet measurements at RHIC?

9

- Different QGP:
  - Temperature/temperature evolution different between LHC and RHIC
- Different probes:
  - Different quark vs. gluon jet mixture
  - Different kinematic range of jets produced







# Jet kinematic reach

10

Projected yields




Projected  $R_{AA}$



- Expect jet measurements out to 70 GeV- overlap with LHC measurements
- High stats for photons ( $\gamma$ -jet measurements) and charged hadrons (fragmentation functions, substructure)

# Jet kinematic reach

11



Calorimeter jet trigger allows for high statistics, high  $p_T$  jet sample

## 3 year run plan projection

Signal	Au+Au 0–10% Counts	p+p Counts
Jets $p_T > 20$ GeV	22 000 000	11 000 000
Jets $p_T > 40$ GeV	65 000	31 000
Direct Photons $p_T > 20$ GeV	47 000	5 800
Direct Photons $p_T > 30$ GeV	2 400	290
Charged Hadrons $p_T > 25$ GeV	4 300	4 100

# Calorimeter jets in sPHENIX

12

- Constituents:  $\Delta\eta \times \Delta\phi = 0.1 \times 0.1$  towers (EMCal + HCals)
- UE subtraction: two iterations, subtract:

$$\frac{d^2 E_T}{d\eta d\phi} = \frac{dE_T}{d\eta} \left( 1 + 2 \sum_n v_n \cos(n(\phi - \Psi_n)) \right)$$

determined  
event-by-event

# Calorimeter jets in sPHENIX

13

- Constituents:  $\Delta\eta \times \Delta\phi = 0.1 \times 0.1$  towers (EMCal + HCals)
- UE subtraction: two iterations, subtract:

$$\frac{d^2 E_T}{d\eta d\phi} = \boxed{\frac{dE_T}{d\eta}} \left( 1 + 2 \sum_n v_n \cos(n(\phi - \Psi_n)) \right)$$

determined  
event-by-event

Average energy  
density, excluding  
regions with jet  
candidates

# Calorimeter jets in sPHENIX

14

- Constituents:  $\Delta\eta \times \Delta\phi = 0.1 \times 0.1$  towers (EMCal + HCals)
- UE subtraction: two iterations, subtract:

$$\frac{d^2 E_T}{d\eta d\phi} = \frac{dE_T}{d\eta} \left( 1 + 2 \sum_n v_n \cos(n(\phi - \Psi_n)) \right)$$

determined  
event-by-event

Average energy  
density, excluding  
regions with jet  
candidates


Flow modulation:  $v_2, v_3, v_4$



# Calorimeter jets in sPHENIX


15

### Jet energy scale (EM scale)



high EM energy scale due to full (EM + hadronic) calorimetry

### Jet energy resolution




low  $p_T$  Au+Au JER high due to UE fluctuations  
• ongoing study to quantify these

# Particle flow jets in sPHENIX

16


- Ongoing work to implement particle flow jets in sPHENIX
- Takes advantage of calorimeter + precision tracking
- Excellent energy response in  $pp$  simulations



# Jet measurements in sPHENIX

17

- Study at RHIC:
  - Path-length dependence of energy loss
  - Mass dependence of energy loss (light vs. heavy flavor jets)
  - Flavor dependence of energy loss (quark vs. gluon jets)
  - How does medium resolve jet substructure?





**Full** characterization of final state

**Different** QGP initial conditions and evolution at RHIC and LHC

**Same** hard process

# Dijet asymmetry

18





- Study path-length dependence of energy loss
- Potential early measurement of jet quenching at RHIC energies

$$A_J = \frac{p_{T,1} - p_{T,2}}{p_{T,1} + p_{T,2}}$$

# Jet $v_2$

19

## Projected yields




- Correlations between energy loss and initial state  $\rightarrow$  path-length dependence of energy loss
- sPHENIX event plane detector (sEPD) allows for measurements of event planes away from jets of interest (see talk by Rosi Reed on Thursday)




# Jet $v_2$

20

Projected yields




Projected  $v_2$



Simultaneous explanation of  $R_{AA}$  and  $v_2$  ongoing "puzzle"

# Photon + jet

21




- High statistics allow for photon + jet measurements
- Photon provides unquenched tag of jet momentum
- Flavor dependence of energy loss

$$x_{J\gamma} = \frac{p_T^{jet}}{p_T^\gamma}$$


# Jet substructure

22

- Fine segmentation of calorimeter + good tracking resolution allows for substructure measurements
- Study how the medium resolves jet substructure




$$z_g = \frac{\min(p_{T,1}, p_{T,2})}{p_{T,1} + p_{T,2}}$$




# b-jets

23

Projected  $R_{AA}$




Projected  $v_2$



- MVTX allows for tagging of heavy-flavor decays
- Study mass dependence of energy loss


# Jets in small systems

24



## □ p+Au data

- Cold nuclear matter effects
- Potential for energy loss in small systems





# Status and timeline

25

- Detector assembly ongoing at BNL
  - ▣ Magnet installed in October
  - ▣ Outer HCal installation nearly complete
  - ▣ Inner HCal and EMCal construction ongoing
- High statistics simulation campaign ongoing
  - ▣ Prep for processing real data + use for performance studies
- Data taking to begin in Feb. 2023



# Summary

26

- sPHENIX detector will provide:
  - ▣ Full coverage electromagnetic and hadronic calorimetry
  - ▣ High precision tracking
  - ▣ Fast readout rate
- Design allows for:
  - ▣ High statistics samples of hard probes (jets, photons, high  $p_T$  charged hadrons)
  - ▣ Full jet reconstruction → complimentary jet measurements to LHC
- Measurements will improve our understanding of small-scale behavior of the QGP

